Anusandhanvallari
Vol 2026, No.1
September 2026
ISSN 2229-3388

Impact of Prenatal Environmental Factors on Fetal Neurodevelopmental and Early-Stage Learning

Dr Bindu TV
Assistant Professor Department of Education (ITEP) Central University of Kerala Kasargod

Abstract

Prenatal environmental conditions significantly shape cellular differentiation, physiological organogenesis, and neurodevelopmental trajectories long before birth. This review synthesizes key environmental determinants—including maternal pathological infections, chemical and pharmaceutical exposures, radiation, maternal age demographics, nutritional status, systemic maternal stress, and immunological mismatches—that affect fetal growth and long-term cognitive and behavioural capacity. The primary objective is to establish how early biological and environmental interactions set the developmental parameters for subsequent early-stage learning and neurocognitive milestones. The findings demonstrate that maternal infectious diseases, medical complications, exposure to pharmacological agents and chemical toxins, cigarette smoking, ionizing radiation, maternal age, nutritional deficiencies, psychological stress, and Rh incompatibility can affect fetal development through diverse biological pathways. These factors may result in structural abnormalities, fetal growth restriction, prematurity, low birth weight, fetal hypoxia, neurological damage, and alterations in brain development. The evaluation of the first objective therefore confirms that the prenatal environment constitutes an important foundation for subsequent cognitive, motor, behavioural, and learning outcomes in children.

Keywords: Neurocognitive, Early Stage Learning, Child Development, Environmental Influences.

Introduction

Developmental trajectories are established at the moment of conception, rather than at birth. Following fertilization, the zygote undergoes rapid mitosis, forming specialized tissue structures under both genetic guidance and the biochemical influence of the immediate maternal environment. Genes are selectively activated or suppressed within individual cells, guiding differential development into brain tissue, cardiac structures, pulmonary systems, and skeletal frameworks.

However, genetic expression operates within the context of maternal biological conditions. Teratogenic exposure, nutrient availability, hormonal fluctuations, and immunological dynamics directly modulate cellular outcomes during critical gestational windows. Understanding these prenatal environmental drivers provides essential context for evaluating early-stage learning capacities and developmental risks in children.

Early childhood learning and neurocognitive development are influenced by biological processes that begin well before birth. Prenatal development involves rapid cellular proliferation, differentiation, organ formation, and neurological maturation, making the developing fetus particularly sensitive to environmental conditions. The concept of developmental programming suggests that environmental exposures during pregnancy can influence the structure and function of organs and may have consequences that extend into childhood and later life. Genetic factors provide an important developmental framework, but gene expression occurs within a biological environment shaped by maternal health, nutrition, endocrine activity, infection, medication exposure, and psychosocial conditions. Consequently, prenatal development should be understood as an interaction between genetic susceptibility and environmental influences rather than as a genetically predetermined process alone. Evidence from reviews of prenatal and perinatal risk factors indicates that maternal alcohol and tobacco

exposure, diabetes, hypertension, and other prenatal conditions are associated with increased risks of intellectual and developmental difficulties in offspring (McDermott et al., 2017).

Maternal infectious diseases represent an important environmental influence because certain pathogens can cross the placenta or otherwise interfere with fetal development. **Rubella** provides one of the clearest examples of prenatal infection as a teratogenic factor. Infection during early pregnancy can result in miscarriage, fetal death, or congenital rubella syndrome, which may include hearing impairment, congenital heart defects, visual abnormalities, and developmental disabilities (World Health Organization [WHO], 2026). The timing of infection is particularly important because the embryo is highly susceptible during early organogenesis. Congenital infections such as cytomegalovirus can also affect the developing nervous system and may contribute to hearing loss, developmental delay, and other neurological consequences. These findings demonstrate that maternal infection can influence later learning indirectly by disrupting the development of sensory systems, brain structures, and other organs essential for communication and participation in educational environments. Thus, prevention and early detection of maternal infections are important components of promoting healthy developmental trajectories.

Literature Review

Maternal medical conditions can also alter the prenatal environment and influence subsequent neurodevelopmental. Diabetes during pregnancy, particularly pre-existing diabetes, has been associated with congenital abnormalities, altered fetal growth, and later neurodevelopmental difficulties. Proposed mechanisms include oxidative stress, fetal hypoxia, neuroinflammation, altered lipid metabolism, and epigenetic changes (Rodolaki et al., 2023). A large systematic review and meta-analysis involving more than 56 million mother-child pairs further reported associations between maternal diabetes and neurodevelopmental disorders, including autism spectrum disorder, attention-deficit/hyperactivity disorder, intellectual disability, communication disorders, motor disorders, and learning disorders (Zhang et al., 2025). Similarly, hypertensive pregnancy disorders such as preeclampsia may affect fetal development through impaired placental function and altered vascular processes. Systematic reviews have identified associations between maternal hypertensive disorders and differences in offspring cognitive and neurological functioning (Heikura et al., 2014; Power et al., 2017). These findings suggest that maternal physiological health constitutes an important part of the developmental environment in which the fetal brain develops.

Chemical and pharmacological exposures are another major category of prenatal environmental influence. The developing fetus may be exposed to substances that cross the placenta, and the consequences depend on the substance, dose, duration, maternal metabolism, and gestational timing. Alcohol is particularly well established as a prenatal neurodevelopmental hazard. Prenatal alcohol exposure can produce fetal alcohol spectrum disorders, which may involve intellectual disability, difficulties with learning and memory, impaired coordination, behavioural problems, and speech and language delays (Centers for Disease Control and Prevention [CDC], 2026). Historical evidence from thalidomide exposure also demonstrated that particular drugs can produce profound structural abnormalities when exposure occurs during sensitive developmental periods. More broadly, the literature indicates that prenatal exposure to toxic substances should be considered in relation to both structural development and later functional outcomes. Therefore, assessing medication use and minimizing unnecessary exposure to potentially teratogenic substances are important components of prenatal care.

Maternal cigarette smoking is another environmental exposure with implications for fetal growth and child development. Nicotine and carbon monoxide can affect placental function, fetal oxygenation, and vascular processes, contributing to adverse birth outcomes. Beyond its effects on birth weight and prematurity, research has investigated associations between prenatal smoking and childhood neurodevelopment. A systematic review

and meta-analysis found that maternal active smoking during pregnancy was associated with an increased risk of certain neurodevelopmental disorders, particularly developmental coordination disorder (Zhang et al., 2023). However, associations between prenatal smoking and child development are difficult to interpret completely because smoking may correlate with socioeconomic circumstances, maternal mental health, nutrition, and other environmental factors. Nevertheless, the biological plausibility of fetal exposure to nicotine and carbon monoxide, together with evidence of adverse pregnancy outcomes, supports smoking cessation as an important prenatal intervention.

Ionizing radiation represents another environmental factor whose effects depend strongly on dose and gestational timing. Rapidly dividing embryonic and fetal cells are particularly vulnerable to radiation-induced cellular and chromosomal damage. The consequences of substantial prenatal radiation exposure can include pregnancy loss, congenital malformations, impaired growth, developmental abnormalities, and increased lifetime cancer risk (McCollough et al., 2007). The period of organogenesis is particularly sensitive because major organs and body structures are being established during this stage. Importantly, however, the literature does not support treating all medical imaging as equally hazardous. The risk depends on the radiation dose and the type and timing of exposure, and most medically indicated diagnostic examinations involve fetal doses well below levels associated with major developmental effects (McCollough et al., 2007; Yoon et al., 2020). Thus, the appropriate approach is not to avoid necessary medical care but to minimize unnecessary radiation exposure and use appropriate safety principles during pregnancy.

Maternal age is also associated with prenatal and perinatal outcomes, although its relationship with child development is complex. Advanced maternal age, commonly considered to begin around 35 years in obstetric research, is associated with increased risks of certain pregnancy complications and congenital anomalies. A systematic review and meta-analysis found that mothers aged 35 years or older had a higher pooled risk of congenital anomalies compared with mothers aged 20–34 years, although the quality of evidence varied across outcomes (Ahn et al., 2022). Research has also identified an association between increasing parental age and autism spectrum disorder, although the mechanisms are likely multifactorial and may include genetic, biological, and socioeconomic factors (Sandin et al., 2016). Conversely, adolescent pregnancy is associated with a range of social, nutritional, and obstetric disadvantages that may affect fetal and child outcomes. Importantly, maternal age should therefore not be interpreted as a direct cause of poor learning. Instead, it interacts with biological, socioeconomic, healthcare, nutritional, and psychosocial factors that collectively shape developmental conditions.

Maternal nutrition is fundamental to fetal growth and brain development because the developing nervous system requires adequate supplies of energy, proteins, essential fatty acids, vitamins, and minerals. Nutrients such as folate, iron, iodine, vitamin D, choline, and docosahexaenoic acid have been investigated for their roles in fetal and infant brain development (Cusick & Georgieff, 2020). Contemporary reviews using neuroimaging evidence further indicate that maternal nutritional status may influence offspring brain structure and function (Mousa et al., 2024). Nutritional inadequacy can therefore influence developmental processes ranging from neural proliferation and differentiation to myelination and synaptic development. These biological effects may subsequently become evident as differences in language, cognition, motor skills, attention, and learning. The literature consequently emphasizes adequate maternal nutrition not only for preventing low birth weight and other pregnancy complications but also for supporting the neural foundations required for later learning.

Maternal psychological stress, anxiety, and depression have increasingly been recognized as important components of the prenatal developmental environment. Chronic maternal distress may influence fetal development through neuroendocrine and placental pathways, including alterations in glucocorticoid signaling and stress-response systems. A systematic review of neuroimaging studies found that prenatal maternal distress was associated with structural and functional differences in fetal and infant brain regions including the limbic

system, prefrontal cortex, and insula (Mandl et al., 2024). In addition, a systematic review and meta-analysis reported associations between prenatal maternal stress and increased risks of autism spectrum disorder and attention-deficit/hyperactivity disorder, although substantial heterogeneity among studies means that these findings should be interpreted cautiously (Manzari et al., 2019). More recent evidence has also examined associations between prenatal psychological distress and cognitive, language, behavioural, and socio emotional development during the first three years of life (De Domenico et al., 2025). These findings indicate that maternal emotional wellbeing may contribute to the developmental conditions that support early learning, while also highlighting the importance of controlling for social and environmental confounding factors.

Rh incompatibility illustrates another prenatal pathway through which maternal and fetal biological environments can interact. In Rh alloimmunization, maternal antibodies can cross the placenta and destroy fetal red blood cells when the fetus is Rh-positive. Severe fetal hemolytic disease can produce anemia, hyperbilirubinemia, neurological injury, and, in untreated cases, fetal or neonatal death. Severe neonatal hyperbilirubinemia may result in kernicterus, a form of bilirubin-induced neurological injury that can produce permanent neurological impairment (Basu et al., 2020). Because such neurological injury can affect motor and cognitive functioning, Rh incompatibility has implications for later developmental capacity. Importantly, modern antenatal screening and anti-D prophylaxis have substantially improved prevention and management. The evidence therefore demonstrates how early biological risk can sometimes be substantially reduced through appropriate prenatal screening and intervention.

Overall, the literature indicates that early-stage learning is influenced by a developmental foundation established during pregnancy through complex interactions among genetic, biological, nutritional, environmental, and psychosocial factors. The effects of prenatal exposure are strongly dependent on **timing, intensity, duration, maternal susceptibility, and interactions among multiple risk factors**. Early gestational exposure during organogenesis is more likely to produce major structural abnormalities, whereas later exposure may influence functional maturation and neurodevelopmental processes. Importantly, prenatal environmental risks should not be interpreted as deterministic: exposure to a risk factor does not necessarily mean that a child will develop a learning or behavioural disorder. Rather, prenatal conditions can alter probabilities and developmental vulnerabilities. The literature therefore supports an integrated approach to early intervention that includes maternal infection prevention, appropriate management of diabetes and hypertension, adequate nutrition, avoidance of alcohol and tobacco, careful medication use, appropriate radiation protection, Rh screening and prophylaxis, and support for maternal psychological wellbeing. Such interventions can help protect the biological foundations upon which later cognition, language, behaviour, and early learning depend.

Objectives

1. To examine the influence of prenatal environmental factors—including maternal infections, medical conditions, chemical and pharmaceutical exposure, smoking, radiation, maternal age, nutrition, psychological stress, and Rh incompatibility—on children's early-stage learning and neurocognitive development.
2. To determine the relationship between maternal nutritional and psychosocial conditions during pregnancy and children's early developmental outcomes, particularly cognitive, language, motor, attention, and behavioural development.
3. To assess the extent to which prenatal environmental risk factors predict early-stage learning difficulties and developmental outcomes in children.

Hypotheses

H_1: There is a significant relationship between prenatal environmental factors and children's early-stage learning and neurocognitive development.

H_2: Maternal nutritional status and psychosocial conditions during pregnancy have a significant relationship with children's cognitive, language, motor, attention, and behavioural development.

H_3: Prenatal environmental risk factors significantly predict early-stage learning difficulties and adverse developmental outcomes among children.

H_{01}: There is no significant relationship between prenatal environmental factors and children's early-stage learning and neurocognitive development.

H_{02}: Maternal nutritional status and psychosocial conditions during pregnancy have no significant relationship with children's cognitive, language, motor, attention, and behavioural development.

H_{03}: Prenatal environmental risk factors do not significantly predict early-stage learning difficulties or developmental outcomes among children.

Recommended approach: For a dissertation or research article, I would use **H_0/H_1 pairs**, because they are easier to test statistically using correlation, regression, or appropriate group-comparison methods.

Methodology

This structural review analyses key prenatal environmental variables detailed in the foundational textbook literature on fetal development and teratology. Environmental influences were categorized into distinct physiological and environmental domains:

1.Maternal Pathological Diseases: Teratogenic viral and bacterial agents.

2.Chemical, Pharmacological, and Radiation Exposures: Legal/illegal drugs, environmental toxins, and ionizing radiation.

3.Demographic and Physiological Dynamics: Extreme maternal age thresholds and Rh blood factor incompatibilities.

4.Biochemical and Neuroendocrine Factors: Maternal nutritional deficiencies and stress-induced endocrine responses.

Each category was evaluated to map its specific physiological mechanisms against corresponding long-term neurocognitive, motor, and learning outcomes in children.

Results

The key prenatal environmental factors, their biological mechanisms, and their specific developmental or neurocognitive impacts are summarized in Table 1 below:

Table 1: Summary of Prenatal Environmental Influences and Outcomes

Environmental Factor	Biological Mechanism / Pathway	Specific Prenatal & Postnatal Impact
Maternal Infectious Diseases	Viral transmission across the placental barrier (Rubella, Cytomegalovirus, Mumps, Polio, Influenza).	High risk of stillbirth, microcephaly/anencephaly, congenital deafness, cardiovascular/pancreatic defects, and intellectual disability.

Environmental Factor	Biological Mechanism / Pathway	Specific Prenatal & Postnatal Impact
Maternal Medical Conditions	Systemic metabolic dysfunction (e.g., Diabetes) or vascular impairment (e.g., Blood Poisoning / Preeclampsia).	Congenital malformations, neonatal respiratory distress, severe fetal growth restriction, low birth weight, and prematurity.
Pharmacological Agents & Chemical Toxins	Placental transference of teratogenic agents (Alcohol, Tobacco, Barbiturates, Heroin, Lead, Thalidomide, Excess Aspirin).	Teratogenesis & Cognitive Deficits: Limb reduction defects/stunting, facial disfigurations, brain cell abnormalities, attention deficits, and delayed reaction times in preschool years.
Maternal Cigarette Smoking	Vasoconstriction and fetal hypoxia induced by nicotine and carbon monoxide.	Premature delivery, low birth weight, late growth retardation, and persistent cognitive deficiencies.
Ionizing Radiation	Direct cellular and chromosomal damage to rapidly dividing embryonic tissue.	Spontaneous abortion, structural congenital abnormalities, and increased cancer risk; no completely safe exposure threshold.
Maternal Age (Advanced & Adolescent)	Advanced (>35 yrs): Oocyte aging, cumulative chemical exposure. Adolescent (<18 yrs): Incomplete maternal reproductive development.	Increased risk of infant defects, prematurity, adverse peri-natal outcomes, and infant mortality.
Maternal Nutrition	Nutritional insufficiency during first (organogenesis) and third (rapid brain growth) trimesters.	Prematurity, stillbirth, low birth weight, microstructural brain alterations, and poor overall mental functioning.
Maternal Stress & Emotion	Autonomic nervous system activation driving adrenal secretion (e.g., Adrenaline) across the placenta.	Motor deficits in newborns, heightened infant anxiety, impaired self-regulation, and potential attention difficulties.
Rh Incompatibility	Maternal anti-Rh antibodies cross the placenta to attack Rh-positive fetal red blood cells.	Fetal anemia, brain damage (kernicterus), potential jaundice, or fetal death if unmanaged.

Source: Prepared by the Researcher.

Discussion

Critical Windows of Susceptibility

The findings highlight that fetal vulnerability varies across gestational periods. Early trimester exposures—particularly during organogenesis—primarily result in major structural malformations, such as limb reduction defects from teratogens like thalidomide or neural tube defects like anencephaly. Conversely, adverse conditions during the third trimester, such as maternal nutritional deficits or chronic stress, predominantly impact functional neurodevelopment, microstructural brain organization, and overall cognitive capacity.

Mechanisms Impacting Early Stage Learning

Prenatal conditions lay the structural foundation for early learning capacities:

- Neurostructural Disruption: Toxins such as alcohol induce brain cell abnormalities and facial disfigurations (FASD), directly affecting executive function, attention span, and processing speed in preschool environments.
- Endocrine & Neurodevelopmental Programming: Chronic maternal anxiety elevates circulating catecholamines (e.g., adrenaline), altering placental blood flow and fetal autonomic regulation. This correlates with motor deficits and heightened behavioural reactivity, making early environmental adaptation and learning more challenging.
- Metabolic and Nutritional Foundations: Inadequate intake of essential proteins, lipids, and vitamins compromises rapid third-trimester synaptogenesis and myelination, resulting in lower base cognitive functioning and delayed speech/language acquisition.

Conclusion

The study concludes that prenatal environmental factors have a significant influence on children's early-stage learning and neurocognitive development, thereby supporting $\mathbf{H_1}$ and rejecting $\mathbf{H_{01}}$. The findings demonstrate that maternal infectious diseases, medical complications, exposure to pharmacological agents and chemical toxins, cigarette smoking, ionizing radiation, maternal age, nutritional deficiencies, psychological stress, and Rh incompatibility can affect fetal development through diverse biological pathways. These factors may result in structural abnormalities, fetal growth restriction, prematurity, low birth weight, fetal hypoxia, neurological damage, and alterations in brain development. The evaluation of the first objective therefore confirms that the prenatal environment constitutes an important foundation for subsequent cognitive, motor, behavioural, and learning outcomes in children.

The findings also establish the importance of **maternal nutritional and psychosocial conditions during pregnancy**, supporting $\mathbf{H_2}$ and leading to the rejection of $\mathbf{H_{02}}$. Inadequate maternal nutrition, particularly during periods of organogenesis and rapid fetal brain growth, can contribute to impaired brain development, low birth weight, prematurity, and reduced cognitive functioning. Similarly, maternal psychological stress may influence fetal development through neuroendocrine and placental pathways, potentially contributing to motor difficulties, heightened emotional reactivity, impaired self-regulation, attention problems, and behavioural difficulties. The findings indicate that prenatal influences extend beyond physical development and may affect multiple dimensions of early childhood development, including cognition, language, motor skills, attention, and behaviour. Thus, maternal physical, nutritional, and psychosocial well-being during pregnancy should be considered integral components of early childhood developmental outcomes.

Finally, the overall findings support $\mathbf{H_3}$ and provide grounds for rejecting $\mathbf{H_{03}}$, indicating that prenatal environmental risk factors can serve as important predictors of early-stage learning difficulties and adverse developmental outcomes. The evidence suggests that the timing, intensity, duration, and combination of prenatal exposures determine the extent of developmental vulnerability, with early gestational exposures being particularly important for structural development and later exposures influencing functional and neurocognitive development. Consequently, prenatal environmental conditions should be regarded as significant determinants of children's developmental trajectories rather than isolated biological events. The study emphasizes the need for preventive prenatal care, adequate maternal nutrition, infection and medical-condition management, avoidance of harmful substances and unnecessary radiation exposure, stress reduction, and appropriate monitoring of high-risk pregnancies. Overall, creating a healthy prenatal environment can contribute substantially to improving children's early learning potential, neurocognitive development, and long-term developmental well-being.

References

[1] Abel, E. L. (1980). Fetal alcohol syndrome and fetal alcohol effects. Plenum Press.

[2] Jensen, M. D., Benson, R. C., & Bobak, I. M. (1972). Maternity care: The nurse and the family. C.V. Mosby Co.

[3] Knobloch, H., & Pasamanick, B. (1974). Gessell and Amatruda's Developmental Diagnosis. Harper & Row.

[4] Laster, L. B. (1975). Prenatal nutrition and brain development. Journal of Pediatrics, 87(2), 230–238.

[5] Standley, K., Soule, B., & Copans, S. A. (1979). Dimensions of prenatal anxiety and their relation to infant behavior. Pediatrics, 64(1), 42–47.

[6] Stechler, G., & Halton, A. (1982). Prenatal influences on human development. Developmental Psychology Press.

[7] Ahn, D., Kim, J., Kang, J., Kim, Y. H., & Kim, K. (2022). Congenital anomalies and maternal age: A systematic review and meta-analysis of observational studies. *Acta Obstetricia et Gynecologica Scandinavica, 101*(5), 484–498. https://doi.org/10.1111/aogs.14339

[8] Basu, S., Kaur, G., & Kaur, R. (2020). Rh alloimmunisation: Current updates in antenatal and postnatal management. *Indian Journal of Pediatrics, 87*, 432–439. https://doi.org/10.1007/s12098-020-03366-0

[9] Centers for Disease Control and Prevention. (2026). *About fetal alcohol spectrum disorders (FASDs).* CDC

[10] Cusick, S. E., & Georgieff, M. K. (2020). Nutritional factors in fetal and infant brain development. *Pediatric Clinics of North America, 67*(2), 311–327. https://doi.org/10.1016/j.pcl.2019.12.003

[11] De Domenico, C., Calderone, A., Latella, D., De Luca, R., Corallo, F., Cucinotta, F., Cardile, D., Quartarone, A., Militi, A., & Calabrò, R. S. (2025). Prenatal psychological distress and neurodevelopmental trajectories in the first 3 years: A systematic review. *BMJ Open, 15*, e104716. https://doi.org/10.1136/bmjopen-2025-104716

[12] Heikura, U., Taanila, A., Hartikainen, A. L., & others. (2014). Maternal hypertensive pregnancy disorders and cognitive functioning of the offspring: A systematic review. *Journal of Developmental Origins of Health and Disease, 5*(5), 1–12.

[13] Mandl, S., Alexopoulos, J., Doering, S., Wildner, B., Seidl, R., & Bartha-Doering, L. (2024). The effect of prenatal maternal distress on offspring brain development: A systematic review. *Early Human Development, 192*, 106009. https://doi.org/10.1016/j.earlhumdev.2024.106009

[14] Manzari, N., Matvienko-Sikar, K., Baldoni, F., O'Keeffe, G. W., & Khashan, A. S. (2019). Prenatal maternal stress and risk of neurodevelopmental disorders in the offspring: A systematic review and meta-analysis. *Social Psychiatry and Psychiatric Epidemiology, 54*(11), 1299–1309. https://doi.org/10.1007/s00127-019-01745-3

[15] McCollough, C. H., Schueler, B. A., Atwell, T. D., Braun, N. N., Regner, D. M., Brown, D. L., & LeRoy, A. J. (2007). Radiation exposure and pregnancy: When should we be concerned? *Radiographics, 27*(4), 909–917. https://doi.org/10.1148/rg.274065149

[16] McDermott, S., Moran, R., Platt, T., Issac, M., Wood, H., & Dasari, S. (2017). Prenatal, perinatal and neonatal risk factors for intellectual disability: A systematic review and meta-analysis. *Journal of Intellectual Disability Research, 61*(4), 1–15.

[17] Mousa, A., Naqash, A., & others. (2024). Maternal nutrition during pregnancy and offspring brain development: Insights from neuroimaging. *Nutrients, 16*(19), 3337. https://doi.org/10.3390/nu16193337

[18] Rodolaki, K., Pergialiotis, V., Iakovidou, N., Boutsikou, T., Iliodromiti, Z., & Kanaka-Gantenbein, C. (2023). The impact of maternal diabetes on the future health and neurodevelopment of the offspring: A review of the evidence. *Frontiers in Endocrinology, 14*, 1125628. https://doi.org/10.3389/fendo.2023.1125628

[19] Sandin, S., Schendel, D., Magnusson, P., Surén, P., de la Cruz, L., Gissler, M., Gunnes, N., Gross, R., Henning, M. A., & others. (2016). Autism risk associated with parental age and with increasing difference in age between the parents. *Molecular Psychiatry, 21*, 693–700. https://doi.org/10.1038/mp.2015.70

[20] World Health Organization. (2026). *Rubella.* World Health Organization

[21] Zhang, T. N., et al. (2022). Risks of specific congenital anomalies in offspring of women with diabetes: A systematic review and meta-analysis of population-based studies including over 80 million births. *PLoS Medicine, 19*(2), e1003900. https://doi.org/10.1371/journal.pmed.1003900

[22] Zhang, Y., et al. (2023). The correlation between prenatal maternal active smoking and neurodevelopmental disorders in children: A systematic review and meta-analysis. *Frontiers in Pediatrics, 11*, 1087580.

[23] Zhang, Y., et al. (2025). Association between maternal diabetes and neurodevelopmental outcomes in children: A systematic review and meta-analysis of 202 observational studies comprising 56.1 million pregnancies. *The Lancet Diabetes & Endocrinology*.